

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 290 047 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: **01.02.95** (51) Int. Cl.⁶: **C07F 9/58, C07F 13/00, C07F 11/00, C07F 15/00, A61K 49/00**
- (21) Application number: **88107346.4**
- (22) Date of filing: **06.05.88**

The file contains technical information submitted after the application was filed and not included in this specification

- (54) **Dipyridoxyl phosphate nuclear magnetic resonance imagery contrast agents.**

- (30) Priority: **08.05.87 US 47614**
- (43) Date of publication of application: **09.11.88 Bulletin 88/45**
- (45) Publication of the grant of the patent: **01.02.95 Bulletin 95/05**
- (84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE
- (56) References cited:
EP-A- 0 053 347
US-A- 3 632 637

BIOORGANIC CHEMISTRY, vol. 15, no. 3, 1987, pages 224-236; D.M. BOWERS-KOMRO et al.: "Single- and Double-Headed Analogs of Pyridoxamine 5'-Phosphate as Probes for Pyridoxamine 5'-Phosphate Utilizing Enzymes"

Int.J.Nucl.Med.Biol.,vol.12,no.5,1985,pages 381 - 386

Inorg.Chem.,27:3530(1988)

- (73) Proprietor: **NYCOMED SALUTAR, INC.**
428 Oakmead Parkway
Sunnyvale,
California 94086 (US)
- (72) Inventor: **Rocklage, Scott M.**
20221 La Paloma Avenue
Saratoga
California 95070 (US)
Inventor: **Quay, Steven C.**
27863 Moody Road
Los Altos Hills
California 94022 (US)
- (74) Representative: **Cockbain, Julian, Dr. et al**
Frank B. Dehn & Co.
Imperial House
15-19, Kingsway
London WC2B 6UZ (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 290 047 B1

Inorg.Chem.,28:477(1989)

G. Elizondo et al. "Hepatobiliary Contrast Agents: Preclinical Efficacy Studies", Abstract published in SMRM-San Francisco, page 795

Description

This invention relates to novel compounds which form highly stable chelates with metal ions and which are useful as metal ion carriers for in vivo medical applications. In particular, this invention is directed to novel dipyridoxyl compounds which form highly stable chelates with polyvalent metal ions, the preparation of the compounds and chelates thereof with polyvalent ions and particularly paramagnetic ions, and the use of the paramagnetic chelates as contrast agents in nuclear magnetic resonance imaging (NMRI).

Background of the Invention

Traditionally, chelates have been used to administer poorly soluble salts in medicine and as antidotes for detoxification in cases of heavy metal or heavy metal isotope poisoning. Chelates have also been used to deliver radioisotopes to areas of the body for imaging and radiation therapy. Most recently, chelates with paramagnetic contrast agents have been reported for use with NMRI.

Paramagnetic metal ions are frequently toxic in the concentrations required for use in NMRI, and introducing them into the body in the form of chelates renders them more physiologically acceptable. This requires that a chelate be able to hold the metal ion tightly in the chelate structure, that is, the formation constant for the chelate must be very large at physiological pH. The paramagnetic metal chelate must also be sufficiently soluble to permit administration of quantities required for imaging in reasonable volumes. Usual routes of administration are orally, intravenously and by enema.

The chelating agent must form a stable chelate with those paramagnetic metals which present a hazard to the body if released during use. Paramagnetic metals which are naturally present in the body are preferred. The chelate forming agent (ligand) must be capable of forming a chelate with a selected paramagnetic material without altering the metal's oxidation state or otherwise reducing its chemical stability.

Since the role of the paramagnetic metal in increasing contrast in NMRI imaging involves reducing the spin-lattice spin relaxation time T_1 and the spin-spin relaxation time T_2 , the chelate structure must hold the metal ion tightly while permitting contact of the metal ion with protons in water molecules.

This invention provides a novel, superior chelating agent and metal complexes therewith which meet the above objectives.

A summary of the history and state of the art of contrast agents for NMRI is presented by Valk, J. et al, BASIC PRINCIPLES OF NUCLEAR MAGNETIC RESONANCE IMAGING. New York: Elsevier, pp 109-114 (1985). The Valk et al publication also describes the imaging equipment and methods for NMRI, and the entire contents of the Valk et al publication are hereby incorporated by reference in their entirety. Chelates with ethylenediaminetetraacetic acid (EDTA) and diethylaminetriaminepentaacetic acid (DTPA) are described. Toxicity problems were reduced by seeking less toxic metal ions such as iron and gadolinium in a complex of gadolinium-DTPA chelate-meglumine. Gadolinium, however, is not naturally present in the body and long term toxicity studies have not been completed. Paramagnetic materials listed in this publication include molecules with unpaired electrons: nitric oxide (NO); nitrogen dioxide (NO_2); and molecular oxygen (O_2). Also included are ions with unpaired electrons, that is ions from the "transition series". Listed ions include Mn^{2+} , Mn^{3+} , Fe^{2+} , Fe^{3+} , Ni^{2+} , Cr^{2+} , Cu^{2+} , the lanthanide series including gadolinium and europium, and nitroxide stable free radicals (NSFR) such as pyrrolidine NSFR and piperidine NSFR. Toxicity problems are indicated to present a major problem with many paramagnetic materials.

Use of alkylenediamine chelates with a variety of paramagnetic ions are described in U.S. Patent 4,647,447. Ferrioxamine-paramagnetic contrast agents are described in U.S. Patent 4,637,929. Manganese(II) is listed as a suitable paramagnetic metal ion for use with polysaccharide derivatives of a variety of chelating compounds including EDTA, DTPA and aminoethyl diphosphonate in PCT application publication no. WO85/05554 (Application No. PCT/GB85/00234). Stable radioactive diagnostics agents containing $^{99\text{m}}\text{Tc}$ chelated with N-pyridoxal-alpha-aminoacids or a pyridoxal salt are disclosed in U.S. Patents 4,313,928, 4,440,739, and 4,489,053.

Taliaferro, C. et al in "New multidentate ligands. 22. N,N'-dipyridoxyethylenediamine-N,N'-diacetic acid: a new chelating ligand for trivalent metal ions", Inorg.Chem. 23:1188-1192 (1984) describe development of N,N'-dipyridoxyethylenediamine-N,N'-diacetic acid (PLED) as a chelating compound for trivalent metal ions. Other chelating compounds described are the Fe(III) chelates of N,N'-ethylenebis-2-(o-hydroxyphenyl)-glycine (EHPG) and N,N'-bis(1-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid (HBED). Properties of chelates of PLED, HBED, EHPG and EDTA with ions of copper, nickel, cobalt, zinc, iron, indium and gallium are compared. Investigation of the structure of PLED is reported by Taliaferro, C. et al, Inorg.Chem. 24:2408-2413 (1985). Green, M. et al, Int.J.Nucl.Med.Biol. 12(5):381-386 (1985) report their evaluation of

PLED as a chelating ligand for the preparation of gallium and indium radiopharmaceuticals, and summarize properties of PLED chelates with Ga(III), In(III), and Fe(III).

Because the compounds of this invention have an aromatic hydroxy group, their value as chelating agents for manganese(II) ions would not be expected; such aromatic hydroxy groups would be expected to react with the manganese(II) ion as an oxidant in the usual way, oxidizing the manganese(II) ion to a higher valence. Frost, et al, *J.Am.Chem.Soc.* 80:530 (1958) report the formation of Mn(II) chelates of EHPG at low pH, but found that attempts to prepare stable manganese(II) complexes with EHPG at higher pH's (above pH 5) was futile as the manganese(II) ion was irreversibly oxidized. This oxidation occurred even under inert atmospheres, and the writers concluded that the oxidation occurred at the expense of the ligand or solvent. Anderegg, G. et al, *Helv.Chim.Acta.* 47:1067 (1964) found the high stability of the Fe(III) chelate of EHPG was due to the high affinity of the Fe(III) ion for the two phenolate groups present in the ionized ligand. L'Eplattenier, F. et al, (*J.Am.Chem.Soc.* 89:837 (1967) describes studies of HBED involving acid titrations of HBED in the presence of a variety of metal ions, including manganese(II). No manganese chelate was isolated, and the manganese products were not characterized. Based on subsequent work by Patch et al, *Inorg.Chem.* 21(8):2972-2977 (1982), it is clear that the manganese (II) ion was oxidized by the phenolic ligand during the titrations of L'Eplattenier et al. Patch et al prepared a Mn(III) complex by reacting Mn(II) salts with EHPG, and concluded the reaction involved the oxidation of the ligand in an irreversible reaction. The ability to maintain Mn(III) in the +3 oxidation state was said to be a unique characteristic of the EHPG molecule. U.S. Patent 3,632,637 describes phenolic chelating agents such as N,N'-di(o-hydroxybenzyl)-ethylenediamine-N,N'-diacetic acid and their use in chelating trivalent and tetravalent metal. These agents are usually stable in the presence of a aromatic hydroxy groups. No use of a compound with an aromatic hydroxy group as a chelating agent for manganese(II) ions is disclosed in these references, confirming the general knowledge about the oxidising properties of the aromatic hydroxy group on manganese compounds, in particular manganese(II) ions.

Dipyridoxyl analogues of pyridoxamine 5'-phosphate having the general structure 1,n-bis(5'-phospho-4'-pyridoxyl)-1,n-diaminoalkane are described in Bowers-Komro et al., *Bioinorganic Chemistry* 15, 224-236 (1987). Such compounds are proposed simply as cleft-depth probes for the catalytic site of enzymes and a chelating function is not reported.

The novel compounds of this invention are shown in Formula I.

30

35

40



45

wherein

R is hydrogen or

50


55



R₁ is hydrogen or



- with the proviso that at least one of R and R₁ is other than hydrogen;
 R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted 1,2-cycloalkylene having from 5 to 8 carbons, or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons; and
 R₄ is hydrogen, hydroxymethyl, alkyl having from 1 to 6 carbons or



- R₅ and R₆ are each, independently, hydroxy, alkoxy having from 1 to 18 carbons, hydroxy-substituted alkoxy having from 1 to 18 carbons, amino or alkylamido having from 1 to 18 carbons; and
 each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons;
 or a salt thereof.

Also included in this invention are the chelates of the compounds of formula I, and salts thereof with metal ions, preferably paramagnetic metal ions having atomic numbers of from 21-29, 42, 44 or 58-70, and optimally manganese (II), and their use as imaging agents.

The novel intermediate compounds from which the compounds of Formula I are prepared are also included within the compounds of this invention.

Fig. 1 shows a structural formula of a species of N,N'-bis(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid, showing the dissociation constants, pK's, as assigned to the protonation sites described in Example 11.

Fig. 2 is a graph showing the relationship between dosage and relaxivity using the Mn(DPDP) compound of this invention based on the data shown in Example 9.

The novel compounds of this invention are shown in Formula I. The pharmaceutically acceptable water-soluble compatible salts of the compounds of Formula I and the chelates thereof are also included within the compounds of this invention.

In Formula I, R₅ and R₆ are preferably each individually hydroxy, alkoxy having from 1 to 8 carbons, 2-hydroxyethoxy, 2,3-dihydroxypropoxy, amino or alkylamido having from 1 to 8 carbons. Optimally, R₅ and R₆ are hydroxy and the salts thereof.


The term "alkyl" and "alkylene", as used herein, include both straight and branch-chained, saturated and unsaturated hydrocarbons. The term "1,2-cycloalkylene" includes both cis and trans cycloalkylene groups and alkyl substituted cycloalkylene groups bonded at the 1,2-positions to respective nitrogen atoms and alkyl substituted derivatives thereof having from 5 to 8 carbons. The term "1,2-arylene" includes phenyl and naphthyl groups bonded at the 1,2-positions to respective nitrogen atoms and alkyl substituted derivatives thereof, having from 6 to 10 carbons.

The compound, N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)ethylenediamine-N,N'-diacetic acid, is referred to hereinafter as DPDP, and the Manganese(II) chelate is referred to hereinafter as Mn(DPDP).


The dicarbonyl compound of Formula I, when R₅ and R₆ are hydroxy and R₃ is ethylene, is DPDP. DPDP has the dissociation constants for the protonation sites shown in Fig. 1. As described in Example 11, at pH of 3 and above, the ligand is anionic and possesses deprotonated metal binding sites, both important criteria for metal chelating agents.

CHELATES


The chelates of this invention are chelates of the compounds of Formula I with metal ions. The chelates can be represented by Formulae II, III or IV.



(II)



(III)



(IV)

In Formulae II, III and IV, Z represents a metal ion and R_3 , R_4 , R_5 and R_6 are the same as described with respect to the compounds of Formula I. The dotted lines in the figure represent the dative bonding between the oxygen and nitrogen atoms and the metal ion. One of the acetyl groups in Formula II is below the plane of the aromatic pyridine rings and the other acetyl group is above the plane of the aromatic pyridine rings, so the metal ion is tightly held within the interior of the chelate salt complex with the

dicarboxy embodiments of this invention. Also included in the chelates of this invention are the pharmaceutically acceptable water-soluble compatible salts, and carboxylic and phosphate group esters with hydroxy-substituted alkanols, alkanols, or alkylamino alcohols, each having from 1 to 18 carbons, of the compounds of Formulas II, III and IV.

5 For use as a medium for NMRI analysis, the central ion of the complex chelate salt must be paramagnetic, and preferably is a divalent or trivalent ion of elements with an atomic number of 21 to 29, 42, 44 and 58 to 70. Suitable ions include chromium(III), manganese (II), iron (III), iron (II), cobalt (II), nickel(II), copper(II), praseodymium(III), neodymium(III), samarium(III), ytterbium(III). Gadolinium(III), terbium(III), dysprosium(III), holmium(III) and erbium(III) are sometimes preferred because of their strong magnetic
10 moments and chemical stability, but because they are not normally present in the body, their long term biological effects are unknown.

With the novel chelate forming compounds of Formula I, chelates of manganese(II) are preferred. Relatively few manganese(II) chelate compounds are known, and only a fraction of these have been characterised i.e., by single crystal X-ray diffraction. Most of the structurally characterised Mn(II) complexes
15 have various mono and bidentate ligands coordinating to the metal center. The Mn(II) complexes of Formulas II, II and III, and the Mn(II) complexes with PLED and the corresponding 1,2-cycloalkylene and 1,2-arylene compounds described in US-A-4935518 issued on June 19th 1990 are the first Mn(II) complexes with a high affinity hexadentate ligand. This configuration provides a more stable and effective form for introducing manganese(II) into the body as a NMRI contrast medium.

20 The manganese(II) complex of Formula II (R_5 and R_6 = hydroxy, R_3 = ethylene), Mn(DPDP), was studied potentiometrically from pH 11.1 to 2.0. The data obtained was analyzed using a model consisting of three one proton steps and one two proton step. Refinement yielded equilibrium constants with calculated e.s.d.'s (log K) of less than 0.02. The log K_f (formation constant) for Mn(DPDP) was calculated to be 14.8 and is identical to that of the manganese chelate of N,N'-bis(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid (HBED) reported by L. Eplathenier, F. et al (supra). The titration data indicates that Mn(DPDP) begins to demetallate as the pH drops below 4.5. At physiological pH, however, the Mn(II) ion is quantitatively bound to the chelating agent, stabilized at the +2 valence.

The chelate forming compounds of this invention are also suitable for forming chelate salts of other metals intended for X-ray diagnosis. In general, these are the elements of an atomic number which is
30 sufficiently high to efficiently absorb X-rays. Diagnostic media containing a physiologically well tolerated chelate salt with central ions of elements with atomic numbers of 57 to 83 are suitable for this purpose. Included in this group are lanthanum(III), gold(III), lead(II), and bismuth(III).

All of the chelates according to this invention useful for NMRI and X-ray analysis are also suitable for use in ultrasonic diagnosis.

35 The elements of the above-listed atomic numbers which form the central ion or ions of the physiologically well tolerated chelate salt, must not be radioactive for the intended use of the diagnostic medium for X-ray diagnosis and NMRI. Radioactive metal ion chelates of the compounds of Formula I are described in U.S. Patent No. 4842845 issued on June 27 1989.

For purposes of clarity, the chelates of this invention will be described hereinafter in terms of
40 paramagnetic ions suitable for use in NMRI analysis. However, this is for purposes of clarity of explanation and not by way of limitation, and chelates of all of the above metal ions are included within the scope of this invention.

If not all of the active hydrogen atoms of the chelates are substituted by the central paramagnetic ion, the solubility of the chelate is increased if the remaining hydrogen atoms are substituted with physiologi-
45 cally biocompatible cations of inorganic and/or organic bases or amino acids. For example, the lithium ion, the potassium ion, the sodium ion and especially the calcium ion are suitable inorganic cations. Suitable cations of organic bases include, for example, ethanolamine, diethanolamine, morpholine, glucamine, N,N-dimethylglucamine, and N-methylglucamine. Lysine, arginine or orithine are suitable as cations of amino acids, as generally are those of other basic naturally occurring acids.

50 The preferred calcium salts have calcium ion to chelating molecule mole ratios of from 0.05 to 1.0, the optimum mole ratios being with the range of from 0.1 to 0.5. At mole ratios of calcium ion to chelate molecule above 1.0, the chelate tends to become insoluble. The soluble calcium salts are most physiologically acceptable since they do not significantly disturb the concentration of free calcium ions in the patient's system.

55 The chelates according to this invention are formed from the chelate forming compounds of Formula I by conventional procedures known in the art. In general, these processes involve dissolving or suspending the metal oxide or metal salt (for example, nitrate, chloride or sulfate) of an element with an atomic number of 21 to 29, 42, 44 or 57 to 83 (for example, oxides or salts of Mn^{+2} , Cr^{+3} , Fe^{+2} , Fe^{+3} , Co^{+3} , Ni^{+2} , Cu^{+2} , Pr^{+3} ,

Nd³⁺, Sm³⁺, Yb³⁺, Gd³⁺, Tb³⁺, Dy³⁺, Ho³⁺, or Er³⁺) in water or a lower alcohol such as methanol, ethanol or isopropanol. To this solution or suspension is added an equimolar amount of the chelating acid in water or a lower alcohol, and the mixture is stirred, if necessary, with heating moderately or to the boiling point, until the reaction is completed. If the chelate salt formed is insoluble in the solvent used, the reaction product is isolated by filtering. If it is soluble, the reaction product is isolated by evaporating the solvent to dryness, for example, by spray drying or lyophilizing.

If acid groups such as the phosphoric acid groups are still present in the resulting chelate, it is advantageous to convert the acidic chelate salt into a neutral chelate salt by reaction with inorganic and/or organic bases or amino acids, which form physiologically biocompatible cations, and to isolate them. This is often unavoidable since the dissociation of the chelate salt is moved toward neutrality to such an extent by a shift in the pH value during the preparation that only in this way is the isolation of homogeneous products or at least their purification made possible. Production is advantageously performed with organic bases or basic amino acids. It can also be advantageous, however, to perform the neutralization by means of inorganic bases (hydroxides, carbonates or bicarbonates) of sodium, potassium or lithium.

To produce the neutral salts, enough of the desired base can be added to the acid chelate salts in an aqueous solution or suspension that the point of neutrality is reached. The resulting solution can then be concentrated to dryness in vacuo. It is often advantageous to precipitate the neutral salts by adding a solvent miscible with water, for example, lower alcohols (methyl, ethyl, isopropyl alcohols, etc.), lower ketones (acetone, etc.), polar ethers (tetrahydrofuran, 1,2-dimethoxyethane, etc.) and thus obtain crystals that isolate easily and purify well. It has been found particularly advantageous to add the desired bases to the reaction mixture even during chelating and thus eliminate a process stage. Other conventional purification procedures such as column chromatography can be used.

Since the chelate salts of Formulas II, III and IV contain a plurality of acid groups, it is possible to produce neutral mixed salts which contain both inorganic and organic physiologically biocompatible cations as counterions. This can be done, for example, by reacting the complexing acids in an aqueous suspension or solution with the oxide or salt of the element supplying the central ion or less than the full amount of an organic base necessary for neutralization, e.g., half, isolating the chelate salt that is formed, purifying it, if desired, and then adding it to the amount of inorganic base necessary for complete neutralization. The sequence of adding the bases can be reversed.

The carboxylic and phosphoric acid groups of the chelating agents can also be neutralized by esterification to prepare carboxylate and phosphate esters. Such esters can be prepared by conventional procedures known in the art, for example, from the corresponding alcohols. Suitable esters include, for example, esters of straight or branch-chained alkanol groups having from 1 to 18 carbons, mono and polyhydric alkyl amino alcohols having from 1 to 18 carbons and preferably from 1 to 6 carbons such as serinol or diethanolamine, and polyhydric alcohols having from 1 to 18 and preferably from 1 to 6 carbons such as ethylene glycol or glycerol.

The diagnostic media for administration is formed using physiologically acceptable media in a manner fully within the skill of the art. For example, the chelate salts, optionally with the addition of pharmaceutically acceptable excipients, are suspended or dissolved in an aqueous medium, and then the solution or suspension is sterilized. Suitable additives include, for example, physiologically biocompatible buffers (as, for example, tromethamine hydrochloride), slight additions of other chelating agents (as for example, diethylenetriaminepentaacetic acid) or, optimally, calcium salts (for example, calcium chloride, calcium ascorbate, calcium gluconate or calcium lactate).

Alternatively, the diagnostic media according to this invention can be produced without isolating the chelate salts. In this case, special care must be taken to perform the chelating so that the salts and salt solutions according to the invention are essentially free of unchelated, potentially toxic metal ions. This can be assured, for example, using color indicators such as xylenol orange to control titrations during the production process. A purification of the isolated salt chelate can also be employed as a final safety measure.

If suspensions of the chelate salts in water or physiological salt solutions are desired for oral administration, a small amount of soluble chelate salt can be mixed with one or more of the inactive ingredients traditionally present in oral solutions such as surfactants, aromatics for flavoring, and the like.

The most preferred mode for administering paramagnetic metal chelates as contrast agents for NMRI analysis is by intravenous administration. Intravenous solutions must be sterile, free from physiologically unacceptable agents, and should be isotonic or iso-osmotic to minimize irritation or other adverse effects upon administration. Suitable vehicles are aqueous vehicles customarily used for administering parenteral solutions such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, and other solutions such as are described in REMINGTON'S

PHARMACEUTICAL SCIENCES. 15th Ed., Easton: Mack Publishing Co. pp 1405-1412 and 1461-1487 (1975) and THE NATIONAL FORMULARY XIV. 14th Ed. Washington: American Pharmaceutical Association (1975), the contents of which are hereby incorporated by reference. The solutions can contain preservatives, antimicrobial agents, buffers and antioxidants conventionally used in parenteral solutions, selecting excipients and other additives which are compatible with the chelates and which will not interfere with the manufacture, storage or use of the products.

The diagnostic media according to this invention can contain from 0.001 to 5.0 moles per liter and preferably from 0.1 to 0.5 moles per liter of the chelate salt.


The chelates of this invention are administered to patients for imaging in amounts which are sufficient to yield the desired contrast. Generally, dosages of from 0.001 to 5.0 mmoles of contrast agent per kilogram of patient body weight are effective to achieve reduction of relaxivity rates. The preferred dosages for most NMRI applications are from 0.02 to 0.5 mmoles of contrast agent per kilogram of patient body weight.

Methods for applying the contrast agents to improve NMRI images, equipment and operating procedures are described by Valk, J. et al, supra. The contrast agents can be used orally and intravenously.

In a novel NMRI application, the contrast agents can be introduced into the cervix, uterus and fallopian tubes. NMR imaging can then be performed to detect causes of infertility such as obstructions or imperfections in the internal surface of the fallopian tubes which might interfere with the movement of the fertilized ovum.


CHELATE FORMING COMPOUNDS

The compounds of Formula I can be formed by reacting the corresponding pyridoxal 5-phosphate (3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridinecarboxyaldehyde) represented by Formula V with a diamine represented by Formula VI.




In the compounds of Formula V and VI, R₃ and R₄ are as defined with respect to Formula I. Pyridoxyl 5-phosphate, pyridoxal, and the other compounds of Formula V, and the alkylenediamine, cycloalkylenediamine and arylene reactants of Formula VI are well known compounds readily available from commercial sources, and they can be readily synthesized by well known procedures fully within the skill of the art.

The reaction of the amino groups of the compounds of Formula VI with the aldehyde group of the compounds of Formula V can be carried out in an alcohol such as methanol at a temperature within the range of from 0 to 60 °C. The diimines formed are represented by Formula VII.



In the compounds of Formula VII, R_3 and R_4 are the same as described with respect to the compounds of Formula I. For the manufacture of compounds wherein R_4 is a phosphonomethyl group, i.e., the 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylideneaminoalkyleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids, 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylideneamino-1,2-cycloalkyleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids, and 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylideneamino-1,2-aryleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids of Formula VI, a diamine of Formula VII is reacted with two molar equivalents of an aldehyde of Formula V having the 5-phosphonomethyl group such as pyridoxyl 5-phosphate. For preparation of compounds of Formula VII wherein R_4 is other than a phosphonomethyl group, the diamine of Formula VI is first reacted with only one molar equivalent of an aldehyde of Formula V having the 5-phosphonomethyl group, and the mono-phosphonomethyl reaction product is reacted with one molar equivalent of a compound of Formula V having the desired R_4 group, such as a 5-hydroxymethyl group, i.e., pyridoxal. The reverse order of reaction can also be used. The reaction products of Formula VII are insoluble in the alcohol and can be isolated by filtration.

The compounds of Formula VII are then hydrogenated by conventional procedures using a palladium or platinum catalyst to yield the diamines of Formula VIII.



30

In the compounds of Formula VIII, R_3 and R_4 are the same as described with respect to the compounds of Formula IV. The 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylaminoalkyleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids, 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylamino-1,2-cycloalkyleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids, 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylamino-1,2-cycloaryleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethyl phosphoric acids, and the monophosphonomethyl compounds of Formula VIII can be left in solution or isolated as crystalline solids.

The compounds of Formula I are prepared by reacting the diamines of Formula VIII with a haloacetic acid such as bromoacetic acid, the molar ratio of the bromoacetic acid to diamine determining whether one or both of the amines are conjugated with the acetic acid groups. The N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)alkylenediamine-N,N'-diacetic acids, N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-1,2-cycloalkylenediamine-N,N'-diacetic acids, N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-1,2-arylenediamine-N,N'-diacetic acids, N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)alkylenediamine-N-acetic acids, N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-1,2-cycloalkylenediamine-N-acetic acids, and N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-1,2-arylenediamine-N-acetic acids of Formula I are then isolated and purified by conventional procedures such as recrystallization or anion exchange chromatography.

The carboxylic acid esters and amides can be formed by conventional procedures reacting the carboxylic acids with alkanols having from 1 to 18 carbons, hydroxy-substituted alkanols having from 1 to 18 carbons, ammonia, and alkylamines having from 1 to 18 carbons.

This invention is further illustrated by the following specific but non-limiting examples. Temperatures are given in degrees centigrade and concentrations as weight percents unless otherwise specified. Procedures which are constructively reduced to practice herein are described in the present tense, and procedures which have been carried out in the laboratory are set forth in the past tense.

EXAMPLE 1

N,N'-bis(pyridoxal-5-phosphate)ethylenediimine

5 A 265.2 gm (1 mole) quantity of pyridoxal-5-phosphate (Chemical Dynamics Corp., South Plainfield, NJ) was slurried in one liter of methanol, and 400 ml of 5 M NaOH was added. When the solution was homogeneous, 34.2 ml of 1,2-diaminoethane (Aldrich Chem. Co.) was added rapidly with vigorous stirring. The imine product sodium N,N'-bis(pyridoxal-5-phosphate)ethylenediimine or sodium 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylideneaminoethyleneiminomethyl)-2-hydroxy-3-methyl-5-
 10 pyridylmethylphosphate was stirred for 1 hr, 400 ml of diethyl ether was added, and the slurry was filtered. The filtrate was washed with 600 ml of ethanol and dried at 60 °C in vacuo. A 290 gm quantity of the bis-imine with a melting point of 215-220 °C (decomposition) was isolated (90% yield, based on the tetra-sodium salt). IR (KBr) pellet: 1630 cm⁻¹ (C=N), ¹H NMR (D₂O, 400 MHz) delta 8.88 (s, -N=CH), 7.54 (s, pyr-H), 4.70 (d, CH₂OP, J_{HP} = 6.3 Hz), 4.06 (s, NCH₂CH₂N), 2.21 (s, pyr-CH₃).

EXAMPLE 2

N,N'-bis(pyridoxal-5-phosphate)alkyldiimines

20 Repeating the procedure of Example 1 but replacing the 1,2-diaminoethane with 1,3-diamino-n-propane, 1,2-diamino-n-propane, 1,2-diaminoisopropane, 1,2-diamino-n-butane, 1,4-diamino-n-butane, 1,3-diamino-n-butane, 1,2-diamino-3-methylpropane yields the corresponding N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-propylene)diimine, N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-propylene)diimine, N,N'-bis(pyridoxal-5-phosphate)-1,2-isopropylenediimine, N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-butylene)diimine, N,N'-bis(pyridoxal-5-phosphate)-1,4-(n-butylene)diimine, N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-butylene)diimine, and N,N'-bis-
 25 (pyridoxal-5-phosphate)-1,2-(3-methyl)propylenediimine.

EXAMPLE 3

N,N'-bis(pyridoxal-5-phosphate)ethylenediamine

The diimine from Example 1 was charged to a 5 liter 3-neck flask fitted with mechanical stirrer, fritted tube bubbler, and a 3-way stopcock. Then 1.5 liters of deionized water was added, followed by 1.5 liters of methanol. The yellow solution formed was stirred while sparging with nitrogen. Then 13 gm of 5% Pt on carbon (Aldrich Chem. Co.) was added, and the apparatus was purged with hydrogen. The reaction was
 35 allowed to proceed for 5 hr with continuous addition of hydrogen. HPLC analysis showed complete reduction to the amine. The reaction mixture was sparged with nitrogen for 15 min and then filtered through Celite. The filtrate was concentrated in vacuo at 60 °C to about 500 ml. The solution, containing N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine or 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)-methylaminoethyleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphoric acid salt was used di-
 40 rectly for the next step. If desired the diamine can be isolated as off-white crystals by the addition of 200 ml of 97% formic acid and allowing the product to crystallize at room temperature overnight. The diamine is isolated by filtration and washed with 2 x 150 ml of cold deionized water. ¹H NMR (D₂O, 400 MHz) delta 7.47 (s, pyr-H), 4.58 (d, CH₂OP, J_{HP} = 6.3 Hz), 3.94 (s, NCH₂CH₂N), 2.88 (s, N-CH₂-pyr), 2.16 (s, pyr-CH₃).

EXAMPLE 4

N,N'-bis(pyridoxal-5-phosphate)alkyldiamines

50 Repeating the procedure of Example 3 but substituting the diimine products of Example 2 for the diimine product of Example 1 yields N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-propylene)diamine, N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-propylene)diamine, N,N'-bis(pyridoxal-5-phosphate)-1,2-isopropylenediamine, N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-butylene)diamine,
 55 N,N'-bis(pyridoxal-5-phosphate)-1,4-(n-butylene)diamine, N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-butylene)diamine, and N,N'-bis(pyridoxal-5-phosphate)-1,2-(3-methyl)propylenediamine.

EXAMPLE 5

DPDP Synthesis

5 The diamine from Example 3 was charged to a two liter 4-neck flask equipped with two addition funnels, pH electrode, thermometer and stir bar. A 100 gm (2.5 mole) quantity of NaOH was dissolved in 200 ml of deionized water, and 130 gm (0.9 mole) of bromoacetic acid (Sigma Chem. Co.) was dissolved in 180 ml of deionized water. Each solution was charged to an addition funnel. Enough NaOH solution was added to the diamine solution to bring the pH to 11. The temperature of the reaction was raised to 42 °C, and
 10 bromoacetic acid and NaOH solution were added concurrently to maintain the pH at 11. The addition was stopped at 45 min, and the progress of the reaction was checked by HPLC. The addition of bromoacetic acid and NaOH was resumed, and the reaction checked at 60 and 75 min. All the bromoacetic acid had been added, and the reaction was complete by HPLC analysis. Approximately 30 ml of the 50% NaOH solution remained in the addition funnel. A 675 gm quantity of cation exchange resin (AMBERLITE IRC-50)
 15 was added, and the mixture was placed in a refrigerator for 14 hr. The pH had dropped to 6.5. The resin was removed by filtration, and the filtrate treated with 260 gm of cation exchange resin (DOWEX 50W-X8). The pH dropped to about 4. The resin was removed by filtration, and the solution was concentrated in vacuo at 60 °C to a viscous oil. The oil was dried in vacuo for 48 hr at 25 °C to yield a resinous solid containing N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)ethylenediamine-N,N'-diacetic acid (DPDP).
 20

EXAMPLE 6

DPDP Purification

25 The resinous solid obtained in Example 5 was dissolved in 600 ml of 88% formic acid, 1.5 liters of methanol followed by 2.2 liters of ethanol was added, and the mixture was cooled to 0 °C for 2 hr. The solvent mixture was decanted from the resulting gum and discarded. The gum was dissolved in about 800 ml of deionized water which was then concentrated in vacuo to about 600-650 ml. Seed crystals were
 30 added, and the solution was allowed to stand at rm temp overnight. The product was isolated by filtration, washed with about 400 ml of cold deionized water, 250 ml of ethanol, and then dried in vacuo to yield 65 gm of DPDP in 85-90% purity by HPLC. The filtrate and washings were retained, concentrated in vacuo to about 350 ml, and the solution refrigerated until column chromatographic purification of the second crop.

The 65 gms of product was then dissolved in 75 ml of 88% formic acid containing 5 ml of deionized
 35 water with gentle heating to about 60 °C. Cold deionized water was added to a total volume of one liter, and the solution was allowed to stand at 25 °C for 16 hr to crystallize. The product was isolated by filtration, washed with 200 ml cold deionized water, and dried in vacuo at 60 °C to yield 55 gms of DPDP in 93-95% purity by HPLC. A second recrystallization, using the same procedure yields 50 gm of DPDP in 96-98% purity by HPLC, mp 174-180 °C with decomposition. Analysis: (Calculated for $C_{22}H_{32}N_4O_{14}P_2$) C, 41.38; H,
 40 5.05; N, 8.77.
 (Found) C, 40.70; H, 5.14; N, 8.61. 1H NMR (D_2O , 400 MHz) δ 7.93 (s, pyr-H), 4.81 (d, CH_2OP , J_{HP} = 6.3 Hz), 4.07 (s, NCH_2CH_2N), 3.35 (s, CH_2COOH), 2.83 (s, $N-CH_2-pyr$), 2.38 (s, $pyr-CH_3$). ^{31}P NMR (D_2O , 161 MHz) δ -1.61 (s, CH_2OP , H_3PO_4 reference).

45 EXAMPLE 7

Sodium-Calcium Salt of Mn(DPDP)

A 4.16 gm (6.25 mmole) portion of DPDP from Example 6 was dissolved in 15 ml of rigorously
 50 degassed water by the addition of 1.0 gm (25 mmoles) of NaOH. A 1.25 gm (6.25 mmole) quantity of manganese dichloride tetrahydrate was added, and the solution immediately turned yellow. After stirring for 30 min, 0.25 gm (6.25 mmole) of solid NaOH was added to bring the pH up to 6.5. Then 0.15 gm (1.0 mmole) of calcium chloride was added, and sufficient degassed water was added to bring the volume of the solution to 25 ml. The clear yellow solution was sterilized by being filtered through a 0.2 micron filter to
 55 yield the sodium-calcium salt of a manganese chelate complex of N,N'-bis-(pyridoxal-5-phosphate)-ethylenediamine-N,N'-diacetic acid or N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-ethylenediamine-N,N'-diacetic acid.

EXAMPLE 8

Relaxivities with Mn(DPDP)

- 5 The relations of protons present in water and plasma exposed to the chelate product of Example 7 was tested by NMR for relaxities, in msec, at 10 MHz, 37°C. The results are shown in Table I.

TABLE I

<u>Relaxivities, msec.</u>				
<u>Molar</u>	<u>T₁</u>	<u>T₂</u>	<u>T₁</u>	<u>T₂</u>
<u>Conc.</u>	<u>(Water)</u>	<u>(Water)</u>	<u>(Plasma)</u>	<u>(Plasma)</u>
0.010	43	41	40	34
0.005	100	86	74	68
0.0025	175		139	124
0.00125	332		240	
0.000625	639		398	
0.000312	1083		624	
0.000156	1470		856	
0.000078			995	
0.000039			1103	

EXAMPLE 9

Organ Distribution of Mn(DPDP) in Rabbits

- 40 Each of four rabbits was injected intravenously with one of the following amounts of the solution obtained in Example 7: 0.01 mmoles/kg, 0.05 mmoles/kg, 0.10 mmoles/kg and 0.20 mmoles/kg. The rabbits were sacrificed 30 min post injection, and the proton relaxation values of selected body organs were measured with NMR, in vitro at 10 MHz. The relaxation rates found are shown in Table II.

TABLE II

Dose (mmol/kg)	<u>Relaxivities, msec.</u>									
	Normal		Observed							
	<u>Values</u>		<u>Values</u>							
			<u>0.01</u>		<u>0.05</u>		<u>0.10</u>		<u>0.20</u>	
<u>Tissue</u>	<u>T₁</u>	<u>T₂</u>	<u>T₁</u>	<u>T₂</u>	<u>T₁</u>	<u>T₂</u>	<u>T₁</u>	<u>T₂</u>	<u>T₁</u>	<u>T₂</u>
Brain	-	-	554	76	496	82	590	90	352	66
Heart	605	70	-	-	353	54	300	51	205	47
Lung	595	112	-	-	575	113	435	61	376	63
Fat	171	154	-	-	200	139	183	115	192	-
Skel.										
Musc.	423	47	-	-	494	47	425	31	232	31
Renal										
Cort.	338	85	298	65	210	57	188	55	143	53
Renal										
Med.	672	149	502	99	223	57	209	48	127	60
Liver	252	64	176	39	76	31	65	23	68	25
Stom.	349	69	-	-	245	41	242	52	271	53
Small										
Int.	352	79	324	72	237	52	218	46	131	46
Large										
Int.	349	77	283	64	365	83	290	57	256	74
Urine	-	-	821	-	150	136	90	75	-	-
Blood	900	-	844	-	613	-	506	-	411	-

The organ distribution data is plotted in Fig. 2 , with the following symbols:

Liver	-l-	Heart	-h-	Cortex	-c-
Medula	-m-	Urine	-u-	Blood	-b-

It shows rapid uptake of the Mn(DPDP) in the heart, liver and kidneys. The liver and kidneys are saturated with a dose of 0.10 mmole/kg while the heart continues to uptake Mn(DPDP) through the dose range studied. The complex of Mn(DPDP) may cross the intake-brain barrier as uptake by the brain was observed at higher doses. In cases where a defect is present in the blood-brain barrier (through disease or trauma), large amounts of the complex of Mn(DPDP) collect in the extravascular space and such defects were observed by NMRI tomography. The same defects are not observable without the use of Mn(DPDP) as a contrast agent.

EXAMPLE 10**Pharmacokinetics with Mn(DPDP)**

Each of seven rabbits was injected intravenously with 0.01 mmol/kg of the solution obtained in Example 7. The rabbits were sacrificed at 0.25, 0.50, 1.0, 2.0, 4, 6 and 24 hours post-injection, and the proton relaxation values of selected body organs were measured with NMR, in vitro, at 10 MHz. The T_1 relaxation rates are shown in Table III.

TABLE III **T_1 Relaxivities, msec**

<u>Time, hr</u>	<u>0.25</u>	<u>0.50</u>	<u>1.0</u>	<u>2</u>	<u>4</u>	<u>6</u>	<u>24</u>	<u>T_1 Normal</u>
								<u>Value</u>
<u>Tissue</u>								
Liver	68	76	48	48	145	209	315	250 \pm 50
Bile	160	46	25	21	<1	14	107	275 \pm 55
Renal								
cortex	202	191	229	192	210	239	328	338 \pm 60
Renal								
medulla	192	231	236	310	340	375	563	672 \pm 100
Heart	231	352	381	507	522	663	660	605 \pm 100

The pharmacokinetic data show rapid uptake and clearance of Mn(DPDP) in the liver, renal cortex, renal medulla and heart. The results indicate clearance of Mn(DPDP) through both the renal and hepatobiliary systems within 6-8 hours post-injection.

EXAMPLE 11**Potentiometric Titrations**

The compound DPDP was studied potentiometrically from pH 11.2 to 2.0. Data sets were collected on a custom-built automatic potentiometric titration apparatus composed of a METROHM 655 DOSIMAT automatic buret, a FISHER ACCUMET pH meter with a CORNING calomel combination electrode, a custom-blown water jacketed titration cell, a BRINKMAN LAUDA K-2/R constant temperature bath and a COM-MODORE 64 computer. The BASIC computer program TITRATOR (Harris, W. et al, J.Am.Chem.Soc. 101:6534 (1979)) runs the apparatus. Data analysis was performed on an IBM-AT computer using the least squares program, BETA (Harris, W. et al, supra), and the data analysis program, HANDNBAR (Harris, W. et al, supra). The titrants were standardized by phenolphthalein titration as follows: KOH was calibrated against potassium hydrogen phthalate (a primary standard), and HCl solutions were calibrated against the KOH standard. The Mn(II)Cl₂ solution was standardized with an EDTA titration using Erichrome Black T as the indicator. All solutions were made from distilled, deionized water that was further purified on a MILLI-Q

cartridge system, degassed, and then kept under an atmosphere of argon which had been scrubbed for CO₂ and O₂. Additions of EDTA and Mn(II) solutions were performed using calibrated GILMOT pipets. The electrode was calibrated in concentration units with degassed solutions of P[H⁺] = 2.291 and 1.078 at 0.1 M ionic strength.

5 The ligand proton titration was performed by adding 28.7 mg (0.045 mmoles) to 54.6 ml of a high pH aqueous solution. It was then titrated to low pH with 0.1009 N HCl.

The metal complex titration was performed by adding 152.2 mg (0.2383 mmoles) to 74.6 ml of a high pH aqueous solution. 2.07 ml of a 0.1152 M Mn(II) solution (0.2386 mmoles) were added. The complex was then titrated to low pH with 1.002 N HCl.

10 The data obtained was analyzed using a model that consisted of eight one-proton steps. The first two protonation equilibria were outside of the range of the titration window afforded by the concentration of titrant and were therefore estimated based on work by Martell and co-workers reported by Taliaferro, C. et al, *Inorg.Chem.* 24:2408-2413 (1985). Refinement of the remaining equilibria yielded constants with calculated e.s.d.'s (log K) of less than 0.02. Assignment of the protonation sites (pK's) was based on work
15 by Martell and co-workers (Taliaferro, et al, *supra*), and is shown in Fig. 1.

At a pH of 3 and above, the ligand is anionic and possesses deprotonated metal binding sites, both important criteria for a metal chelating agent.

EXAMPLE 12

20

N,N'-bis-(pyridoxal-5-phosphate)alkylenediamine-N,N'-diacetic acids

Repeating the procedure of Examples 5 and 6 but replacing the diamine of Example 3 with the products of Example 4 yields

25 N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-propylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-propylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-isopropylene-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-butylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,4-(n-butylene)-N,N'-diacetic acid,
30 N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-butylene)-N,N'-diacetic acid, and
N,N'-bis(pyridoxal-5-phosphate)-1,2-(3-methylene)propyl-N,N'-diacetic acid.

EXAMPLE 13

35 DPDP Chelates

Repeating the procedure of Example 7 but replacing manganese dichloride tetrahydrate with equimolar amounts of the soluble chlorides of Cr⁺³, Fe⁺², Fe⁺³, Co⁺³, Ni⁺², Cu⁺², Pr⁺³, Nd⁺³, Sm⁺³, Yb⁺³, Gd⁺³, Tb⁺³, Dy⁺³, Ho⁺³, or Er⁺³ yields the corresponding sodium salts of the respective metal ion chelates of N,N'-bis-
40 (pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid. The procedure can be repeated replacing the metal chloride salts with soluble nitrate or sulfate salts.

EXAMPLE 14

45 Other Chelates

Repeating the procedure of Example 7 but replacing N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid with equimolar amounts of the chelate forming compounds produced in accordance with Example 10, and replacing manganese dichloride tetrahydrate with equimolar amounts of the soluble
50 chlorides, carbonates or nitrates of Cr⁺³, Fe⁺², Fe⁺³, Co⁺³, Ni⁺², Cu⁺², Pr⁺³, Nd⁺³, Sm⁺³, Yb⁺³, Gd⁺³, Tb⁺³, Dy⁺³, Ho⁺³, or Er⁺³ yields the sodium-calcium salts of the respective metal ion chelates of

N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-propylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-propylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-isopropylene-N,N'-diacetic acid,
55 N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-butylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,4-(n-butylene)-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-butylene)-N,N'-diacetic acid, and
N,N'-bis(pyridoxal-5-phosphate)-1,2-(3-methylene)propyl-N,N'-diacetic acid.

EXAMPLE 15

N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediimine

- 5 A 26.5 gm quantity (0.1 mole) of pyridoxal-5-phosphate was dissolved in 300 ml of methanol, and 38 ml of 5 N NaOH was added. Then 5.71 gm (0.05 mole) of trans-1,2-diaminocyclohexane was added with stirring, and the volume of the solution was reduced to 200 ml in vacuo. After cooling to 0 °C, the yellow imine was isolated by filtration, washed with diethyl ether, and dried in vacuo to yield 17 gm of sodium N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediimine or sodium 5-(N-(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridyl)methylideneamino-trans-1,2-cyclohexyleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphate (49% yield, melting point 200-205 °C with decomposition).

EXAMPLE 16

- 15 Other N,N'-bis(pyridoxal-5-phosphate)-1,2-cyclo(alkylene or arylene)diimines

Repeating the procedure of Example 15 but replacing the trans-1,2-diaminocyclohexane with trans-1,2-diaminocyclopentane, trans-1,2-diaminocycloheptane, trans-1,2-diaminocyclooctane, cis-1,2-diaminocyclohexane, trans-1,3-diaminocyclohexane, trans-1,4-diaminocyclohexane, o-aminoaniline and cis-1,4-diaminocyclohexane yields the corresponding

20 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclopentylenediimine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cycloheptylenediimine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclooctylenediimine,
N,N'-bis(pyridoxal-5-phosphate)-cis-1,2-cyclohexylenediimine,
25 N,N'-bis(pyridoxal-5-phosphate)-trans-1,3-cyclohexylenediimine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,4-cyclohexylenediimine,
N,N'-bis(pyridoxal-5-phosphate)-1,2-phenylenediimine, and
N,N'-bis(pyridoxal-5-phosphate)-cis-1,4-cyclohexylenediimine.

EXAMPLE 17

N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediamine

- A 14 gm (0.02 mole) portion of the diimine product of Example 15 was dissolved in 200 ml of 1:1 water: methanol. The resulting solution was sparged with argon, and 1.0 gm of 5% platinum on carbon was added. The system was flushed with hydrogen. The hydrogen pressure was increased to 50 psig for 16 hr at 25 °C. The reaction product was filtered through CELITE, and the resulting solution of N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexyldiamine or sodium 5-(N-(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridyl)methylideneamino-trans-1,2-cyclohexyliminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphate was concentrated in vacuo to about 20 ml and cooled to 0 °C to induce crystallization. The product was isolated by
- 40 filtration, washed with cold H₂O and dried in vacuo. ¹H NMR (D₂O, 400 MHz) delta 7.45 (s, pyr-H), 4.53 (d, CH₂OP, J_{HP} = 4.9 Hz), 3.83 (dd, N-CH₂-pyr), 2.72 (br s, cyclo-(CH₂)₄(CH)₂(NH)₂-), 1.88 (s, pyr-CH₃), 1.83-1.08 (3 br s, cyclo-(CH₂)₄(CH)₂(NH)₂-).

EXAMPLE 18

N,N'-bis(pyridoxal-5-phosphate)-1,2-cyclo(alkylene or arylene)diamines

- Repeating the procedure of Example 17 but replacing the diimine product of Example 15 with the diimine products prepared in accordance with the procedure of Example 16 yields the corresponding diamines:
- 50 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclopentylenediamine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cycloheptylenediamine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclooctylenediamine,
55 N,N'-bis(pyridoxal-5-phosphate)-cis-1,2-cyclohexylenediamine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,3-cyclohexylenediamine,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,4-cyclohexylenediamine,
N,N'-bis(pyridoxal-5-phosphate)-1,2-phenylenediamine, and

N,N'-bis(pyridoxal-5-phosphate)-cis-1,4-cyclohexylenediamine.

EXAMPLE 19

5 N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediamine-N,N'-diacetic acid

The diamine from Example 17 was charged to a one liter 3-neck flask, and the pH was adjusted to 11 with 5 N NaOH. Then 5.6 gm (0.04 mole) of bromoacetic acid was dissolved in 10 ml of water and added dropwise to the stirred diamine solution while maintaining the pH at 11. The reaction was warmed to 50 °C and stirred for 16 hr. 50 gm of weakly acidic cation exchange resin (AMBERLITE IRC-50) was added, and the pH dropped to 6.7. The resin was removed by filtration, and 15 gm of cation exchange resin (DOWEX 50W-X8) was added. The pH dropped to 3.8.

The solution was filtered, and all of the solvent was evaporated from the filtrate to yield a foamy solid. The solid was dissolved in 30 ml of 88% formic acid, and the product was precipitated by the addition of 150 ml of methanol followed by 150 ml of ethanol. The solvent mixture was decanted from the gummy solid and discarded. The solid was dissolved in a minimum amount of deionized water (about 100 ml), and the product was allowed to stand overnight at 25 °C. The product was isolated by filtration, washed with 50 ml of cold water, 25 ml of ethanol and then dried in vacuo to yield the product. The compound was recrystallized by the same procedure to yield N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediamine-N,N'-diacetic acid or N,N'-bis(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)-trans-1,2-cyclohexylenediamine-N,N'-diacetic acid (DPCP) with a melting point (decomposition) of 221-226 °C. ¹H NMR (D₂O, 400 MHz) delta 7.53 (s, pyr-H), 4.58 (d, CH₂OP, J_{HP} = 5.9 Hz), 3.89 (dd, N-CH₂-pyr), 3.31 (s, CH₂COOH), 2.78 (br s, cyclo-(CH₂)₄(CH)₂(NH)₂-), 1.93 (s, pyr-CH₃), 1.90-1.15 (3 br s, cyclo-(CH₂)₄(CH)₂-(NH)₂-).

EXAMPLE 20

N,N'-bis-(pyridoxal-5-phosphate)-cyclo(alkyleneand arylene)diamine-N,N'-diacetic acids

Repeating the procedure of Example 19 but replacing the diamine of Example 17 with the diamines of Example 18 yields the corresponding:

N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclopentylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cycloheptylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclooctylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-cis-1,2-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,3-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,4-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-phenylenediamine-N,N'-diacetic acid, and
N,N'-bis(pyridoxal-5-phosphate)-cis-1,4-cyclohexylenediamine-N,N'-diacetic acid.

EXAMPLE 21

Chelates

Repeating the procedure of Example 7 but replacing N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid with equimolar amounts of the products of chelate forming compounds produced in accordance with Examples 19 and 20 and using equimolar amounts of the soluble chlorides, carbonates or nitrates of Mn⁺², Cr⁺³, Fe⁺², Fe⁺³, Co⁺³, Ni⁺³, Cu⁺², Pr⁺³, Nd⁺³, Sm⁺³, Yb⁺³, Gd⁺³, Tb⁺³, Dy⁺³, Ho⁺³, or Er⁺³ yields the sodium-calcium salts of the respective metal ion chelates of

N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclopentylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cycloheptylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclooctylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-cis-1,2-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,3-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-trans-1,4-cyclohexylenediamine-N,N'-diacetic acid,
N,N'-bis(pyridoxal-5-phosphate)-1,2-phenylenediamine-N,N'-diacetic acid, and
N,N'-bis(pyridoxal-5-phosphate)-cis-1,4-cyclohexylenediamine-N,N'-diacetic acid.

EXAMPLE 22

Pyridoxal-5-phosphate (N-methylethanolamine)monoester

- 5 2.04 gm (0.01 mole) of pyridoxal hydrochloride is dissolved in 50 ml of dry THF containing 0.05 gm (0.02 mole) of sodium hydride with stirring. When gas evolution had ceased (about 15 min), 1.71 gm (0.01 mole) of benzyl bromide is added, and after stirring overnight, the solution is brought to dryness in vacuo. The sticky solid is suspended in 50 ml of dry methylene chloride and following addition of 3.0 gm (0.03 mole) of triethylamine, the slurry is cooled to 0 °C. 1.38 gm (0.01 mole) of 2-chloro-3-methyl-1-oxa-3-aza-2-phosphacyclopentane (prepared by the method of Jones, et al, *J.Chem.Soc. Perkin trans I.* p 199 (1985)) is added with vigorous stirring. The suspension is stirred for 1 hr at rm temp, and then 100 ml of water is added. The methylene chloride layer is separated, dried over MgSO₄, and the solvent removed in vacuo. Addition of diethyl ether yields the intermediate product as a hygroscopic white solid (1.8 gm, 50% yield). The intermediate is oxidized with excess dinitrogen tetroxide in methylene chloride at -78 °C, and then is treated with aqueous HCl in THF under reflux to give the (N-methylethanolamine)monoester of pyridoxal-5-phosphate in an overall yield of 40%.

EXAMPLE 23

- 20 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)ethylenediimine

Repeating the procedure of Example 1 but replacing pyridoxal-5-phosphate with the product of Example 22 yields N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)ethylenediimine or sodium 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)methylideneaminoethyleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphoric acid, N-methylethanolamine ester.

EXAMPLE 24

Other Monoester Diimines

- 30 Repeating the procedure of Example 23 with 1,3-diamino-n-propane, 1,2-diamino-n-propane, 1,2-diaminoisopropane, 1,2-diamino-n-butane, 1,4-diamino-n-butane, 1,3-diamino-n-butane, and 1,2-diamino-3-methylpropane yields the corresponding N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-propylene)diimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-propylene)diimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-isopropylenediimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-butylene)diimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,4-(n-butylene)diimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-butylene)diimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(3-methyl)propylenediimine. Repeating the procedure of Example 23 with trans-1,2-diaminocyclohexane, trans-1,2-diaminocyclopentane, trans-1,2-diaminocycloheptane, trans-1,2-diaminocyclooctane, o-aminoaniline, and cis-1,2-diaminocyclohexane, yields the corresponding N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclohexylenediimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclopentylenediimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cycloheptylenediimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester) trans-1,2-cyclooctylenediimine, N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-phenylenediimine, and N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-cis-1,2-cyclohexylenediimine.

EXAMPLE 25

N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)ethylenediamine

- 55 Repeating the procedure of Example 3 but substituting the diimine product of Example 23 for the diimine product of Example 1 yields N,N'-bis(pyridoxal-5-phosphate(N-methyl-ethanolamine)monoester)-ethylenediamine or 5-(N-(3-hydroxy-2-methyl-5-phosponomethyl-4-pyridyl)-methylaminoethyleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphoric acid, N-methyl-

ethanolamine ester.

EXAMPLE 26

5 Other Monoester Diamines

Repeating the procedure of Example 25 but substituting the diimine products of Example 24 for the diimine product of Example 23 yields

- 10 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-propylene)diamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-propylene)diamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-isopropylenediamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-butylene)diamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,4-(n-butylene)diamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-butylene)diamine,
 15 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(3-methyl)propylenediamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclohexylenediamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclopentylenediamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cycloheptylenediamine,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclooctylenediamine,
 20 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-phenylenediamine, and
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-cis-1,2-cyclohexylenediamine.

EXAMPLE 27

25 DPDP-phosphate monoester

Repeating the procedure of Examples 5 and 6 but replacing the diamine of Example 3 with the product of Example 25 yields N,N'-bis(pyridoxal-5-phosphate(N-methyl-ethanolamine)monoester)ethylenediamine-
 N,N'-diacetic acid, sodium salt or N-methylethanolamine phosphate ester of 5-(N-(3-hydroxy-2-methyl-5-
 30 phosponomethyl-4-pyridyl)methylaminoethyleneaminomethyl)-2-hydroxy-3-methyl-5-
 pyridylmethylphosphoric acid, sodium salt.

EXAMPLE 28

35 Other Diamine-N,N'-diacetic Acid Phosphate Monoesters

Repeating the procedure of Example 27 but replacing the products of Example 26 for the product of Example 25 yields

- 40 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-propylene)diamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-propylene)diamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-isopropylenediamine-N,N'-diacetic acid salt,
 45 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(n-butylene)diamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,4-(n-butylene)diamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,3-(n-butylene)diamine-N,N'-diacetic acid salt,
 50 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-(3-methyl)propylenediamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclohexylenediamine-N,N'-diacetic acid salt,
 55 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclopentylenediamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cycloheptylenediamine-N,N'-diacetic acid salt,

N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-trans-1,2-cyclooctylenediamine-N,N'-diacetic acid salt,
 N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-1,2-phenylenediamine-N,N'-diacetic acid salt, and N,N'-bis(pyridoxal-5-phosphate(N-methylethanolamine)monoester)-cis-1,2-cyclohexylenediamine-
 5 N,N'-diacetic acid salt.

EXAMPLE 29

Chelates

10 Repeating the procedure of Example 7 but replacing N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid with an equimolar amount of the chelate forming compound produced in accordance with Example 27, yields the sodium-calcium Mn(II) chelate of N,N'-bis(pyridoxal-5-phosphate-(N-methyl-ethanolamine)monoester)ethylenediamine-N,N'-diacetic acid.

15 Repeating this procedure, replacing manganese dichloride tetrahydrate with equimolar amounts of the soluble chlorides, carbonates or nitrates of Cr⁺³, Fe⁺², Fe⁺³, Co⁺³, Ni⁺², Cu⁺², Pr⁺³, Nd⁺³, Sm⁺³, Yb⁺³, Gd⁺³, Tb⁺³, Dy⁺³, Ho⁺³, or Er⁺³ yields the sodium-calcium salts of the respective metal ion chelates of N,N'-bis-(pyridoxal-5-phosphate-(N-methyl-ethanolamine)monoester)ethylenediamine-N,N'-diacetic acid.

EXAMPLE 30

Other Chelates

25 Repeating the procedures of Example 29 but replacing the products of Example 28 for the product of Example 27 yields the Mn⁺², Cr⁺³, Fe⁺², Fe⁺³, Co⁺³, Ni⁺², Cu⁺², Pr⁺³, Nd⁺³, Sm⁺³, Yb⁺³, Gd⁺³, Tb⁺³, Dy⁺³, Ho⁺³, or Er⁺³ ion chelates of the sodium-calcium salts of the diacetic acid chelating agents of Example 28.

EXAMPLE 31

DPDP-(mono)acetic Acid Analog Synthesis

10 10 gms (0.017 mole) of the diamine from Example 3 was dissolved in 25 ml of 1:1 water/methanol and charged to a 250 ml 4-neck flask equipped with two addition funnels, pH electrode, thermometer and stir bar. 1.4 gms (0.035 mole) of NaOH and 2.4 gm (0.017 mole) of bromoacetic acid were each dissolved in 10
 35 ml of deionized water and charged to the two addition funnels. Sufficient NaOH was added to the stirring diamine solution to bring the pH to about 11, which raised the temperature to about 40 °C. The temperature was maintained at 40 °C, and bromoacetic acid and NaOH were added concurrently to maintain the pH at 11 over the course of 3 hr. The reaction was monitored by HPLC. Dowex 50W-X8 resin was added to lower the pH from 11.1 to 3.1, the solution was filtered, and resin was washed with 100 ml of deionized water. The
 40 pH of the filtrate was about 3.3. 5 ml of 97% formic acid was added, and the pH dropped to 3.0. Then 10 ml of isopropyl alcohol was added with a few seed crystals, the product stirred overnight at 30 to 40 °C, and then allowed to cool to 25 °C. The crude product was collected by filtration and washed with deionized water. The crude product was then dried at 50 °C in vacuo to yield 3 gms of product (30% yield). The product can be recrystallized from a formic acid/water mixture to yield 2.4 gms in 96-98% purity by HPLC
 45 to yield N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N-acetic acid.

EXAMPLE 32

Other (Mono)acetic Acids

50 Repeating the procedure of Example 31 with the products of Examples 4, 17 and 18 yields the corresponding

N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-propylene)diamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-propylene)diamine-N-acetic acid,
 55 N,N'-bis(pyridoxal-5-phosphate)-1,2-isopropylendiamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-1,2-(n-butylene)diamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-1,4-(n-butylene)diamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-1,3-(n-butylene)diamine-N-acetic acid,

N,N'-bis(pyridoxal-5-phosphate)-1,2-(3-methyl)propylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclohexylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclopentylene-diamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cycloheptylene-diamine-N-acetic acid,
 5 N,N'-bis(pyridoxal-5-phosphate)-trans-1,2-cyclooctylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-cis-1,2-cyclohexylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-trans-1,3-cyclohexylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-trans-1,4-cyclohexylenediamine-N-acetic acid,
 N,N'-bis(pyridoxal-5-phosphate)-1,2-phenylenediamine-N-acetic acid, and
 10 N,N'-bis(pyridoxal-5-phosphate)-cis-1,4-cyclohexylenediamine-N-acetic acid.

EXAMPLE 33

Other Chelates

15

Repeating the procedures of Example 29 but replacing the products of Example 32 for the product of Example 27 yields the Mn^{+2} , Cr^{+3} , Fe^{+2} , Fe^{+3} , Co^{+3} , Ni^{+2} , Cu^{+2} , Pr^{+3} , Nd^{+3} , Sm^{+3} , Yb^{+3} , Gd^{+3} , Tb^{+3} , Dy^{+3} , Ho^{+3} , or Er^{+3} ion chelates of the sodium-calcium salts of the monoacetic acid chelating agents of Example 32.

20

EXAMPLE 34

N-pyridoxal-N'-(pyridoxal-5-phosphate)ethylenediimine

25

A 25 gm (0.123 mole) quantity of pyridoxal hydrochloride is slurried in 100 ml of methanol, and 4.88 gm (0.123 mole) of NaOH is added. When the solution is homogeneous, it is added dropwise to 7.5 gm of 1,2-diaminoethane in 100 ml of methanol with stirring. After 60 min, a methanol solution containing 32.7 gm (0.123 mole) of pyridoxal-5-phosphate and 4.88 gm (0.123 mole) of NaOH is added with vigorous stirring. The unsymmetrical imine product, 5-(N-(3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridyl)-methylideneaminoethyleneiminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphoric acid or
 30 N-pyridoxal-N'-(pyridoxal-5-phosphate)-ethylenediimine, is stirred for 1 hr, and the product is isolated by filtration. The diimine is washed with methanol (2 x 50 ml) and diethyl ether (2 x 50 ml), and dried in vacuo.

EXAMPLE 35

35

N-pyridoxal-N'-(pyridoxal-5-phosphate)ethylenediamine

Repeating the procedure of Example 3 but substituting the diimine product of Example 34 for the diimine product of Example 1 yields the corresponding diamine product,

40

5-(N-(3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridyl)methylaminoethyleneaminomethyl)-2-hydroxy-3-methyl-5-pyridylmethylphosphoric acid or
 N-pyridoxal-N'-(pyridoxal-5-phosphate)-ethylenediamine.

EXAMPLE 36

45

DPMP Synthesis

Repeating the procedures of Example 5 and 6 with the product of Example 35 yields the corresponding N-pyridoxal-N'-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or

50

N-(3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridylmethyl)-N'-(3-hydroxy-2-methyl-5-phosphonomethyl-4-pyridylmethyl)ethylenediamine-N,N'-diacetic acid (DPMP).

EXAMPLE 37

55

Other Chelates


Repeating the procedures of Example 29 but replacing the product of Example 36 for the product of Example 27 yields the Mn^{+2} , Cr^{+3} , Fe^{+2} , Fe^{+3} , Co^{+3} , Ni^{+2} , Cu^{+2} , Pr^{+3} , Nd^{+3} , Sm^{+3} , Yb^{+3} , Gd^{+3} , Tb^{+3} , Dy^{+3} ,

Ho⁺³, or Er⁺³ ion chelates of the sodium-calcium salts of N-pyridoxal-N'-(pyridoxal-5-phosphate)-ethylenediamine-N,N'-diacetic acid.

Claims

5 **Claims for the following Contracting States : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL, SE**

1. A compound of formula I



(wherein

25 R is hydrogen or



R₁ is hydrogen or



40

with the proviso that at least one of R and R₁ is other than hydrogen;

R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted

1,2-cycloalkylene having from 5 to 8 carbons, or optionally alkyl substituted 1,2-arylene having from 6 to 10 carbons; and


45 R₄ is hydrogen, hydroxymethyl, alkyl having from 1 to 6 carbons or




55 R₅ and R₆ are each, independently, hydroxy, alkoxy having from 1 to 18 carbons, hydroxy-substituted alkoxy having from 1 to 18 carbons, amino or alkylamido having from 1 to 18 carbons; and each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.

2. A compound as claimed in Claim 1 wherein for formula I each R_7 is hydrogen.

3. A compound as claimed in Claim 1 wherein for formula I R is



and R_1 is



4. A compound as claimed in Claim 3 wherein for formula I R_5 and R_6 are each, independently, hydroxy, alkoxy having from 1 to 8 carbons, amino or alkylamido having from 1 to 8 carbons.

5. A compound as claimed in Claim 4 wherein for formula I R_5 and R_6 are hydroxy.

6. A compound as claimed in Claim 5 being N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.

7. A chelate of a chelating compound as claimed in any one of claims 1 to 6 with a metal ion having an atomic number of from 21 to 29, 42, 44 or 58-70, or a salt thereof.


8. A chelate as claimed in Claim 7 wherein the metal ion is selected from chromium(III), manganese(II), iron(III), iron(II), cobalt(II), nickel(II), copper(II), praseodymium(III), neodymium(III), samarium(III), ytterbium(III), gadolinium(III), terbium(III), dysprosium(III), holmium(III) and erbium(III).

9. A chelate as claimed in Claim 8 being a manganese (II) chelate of N,N'-bis-(pyridoxal-5-phosphate)-ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.

10. A calcium salt of a chelate as claimed in any one of Claims 7 to 9.

11. A calcium salt as claimed in Claim 10 wherein the molar ratio of calcium to the chelating compound of formula I is from 0.05 to 1.0.


12. A chelating compound intermediate of formula (VII)



15 wherein


R_3 is alkylene having from 1 to 8 carbons, optionally alkyl-substituted cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R_4 is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or



each R_7 independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.


30 13. A chelating compound intermediate of formula (VIII)



45 wherein

R_3 is optionally alkyl-substituted 1,2-cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R_4 is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or




each R_7 independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.


14. A magnetic resonance imaging contrast medium composition comprising a chelate or a calcium salt thereof as claimed in any one of claims 7 to 11 together with a pharmaceutically acceptable, compatible excipient.
15. Use of a chelate or a calcium salt thereof as defined in any one of claims 7 to 11 for the preparation of a magnetic resonance imaging contrast medium composition.
16. A method of imaging comprising administering to a subject a chelate or a calcium salt thereof as claimed in any one of claims 7 to 11 and generating an image of at least a part of said subject.

Claims for the following Contracting State : ES


1. A process for preparing a compound of formula I




(wherein
R is hydrogen or




R₁ is hydrogen or



with the proviso that at least one of R and R₁ is other than hydrogen;
R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted
1,2-cycloalkylene having from 5 to 8 carbons, or optionally alkyl substituted 1,2-arylene having from 6
to 10 carbons; and
R₄ is hydrogen, hydroxymethyl, alkyl having from 1 to 6 carbons or




R_5 and R_6 are each, independently, hydroxy, alkoxy having from 1 to 18 carbons, hydroxy-substituted alkoxy having from 1 to 18 carbons, amino or alkylamido having from 1 to 18 carbons; and each R_7 independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof, said process comprising reacting a compound of formula VIII




(wherein R_3 , R_4 and R_7 are as hereinbefore defined) with a haloacetic acid, and optionally converting the compound obtained into a said salt.

- 20
2. A process as claimed in Claim 1 wherein in said compound of formula VIII each R_7 is hydrogen.
 3. A process as claimed in Claim 1 wherein in said compound of formula I R is




30 and R_1 is



- 40
4. A process as claimed in Claim 3 wherein in said compound of formula I R_5 and R_6 are each, independently, hydroxy, alkoxy having from 1 to 8 carbons, amino or alkylamido having from 1 to 8 carbons.
 5. A process as claimed in Claim 4 wherein in said compound of formula I R_5 and R_6 are hydroxy.
 - 45
 6. A process as claimed in Claim 5 wherein said compound of formula I is N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.
 - 50
 7. A process for preparing a chelate of a chelating compound as claimed in any one of claims 1 to 6 or a salt thereof, said process comprising metallating said chelating compound or salt thereof with a metal ion having an atomic number of from 21 to 29, 42, 44 or 58-70.
 - 55
 8. A process as claimed in Claim 7 wherein the metal ion is selected from chromium(III), manganese(II), iron(III), iron(II), cobalt(II), nickel(II), copper(II), praseodymium(III), neodymium(III), samarium(III), ytterbium(III), gadolinium(III), terbium(III), dysprosium(III), holmium(III) and erbium(III).

9. A process as claimed in Claim 8 for preparing a manganese(II) chelate of N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.
10. A process as claimed in any one of claims 7 to 9 for preparing a calcium salt of a chelate as defined in any one of Claims 7 to 9.
11. A process as claimed in Claim 10 wherein in said calcium salt the molar ratio of calcium to the chelating compound of formula I is from 0.05 to 1.0.
12. A process for preparing a chelating compound intermediate of formula (VII)




(VII)


wherein

R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R₄ is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or



each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof, said process comprising reacting a compound of formula V




(V)

(wherein R₄ is as hereinbefore defined) with a compound of formula VI



(wherein R₃ is as hereinbefore defined) and optionally converting the compound obtained into a said salt.


13. A process for preparing a chelating compound intermediate of formula (VIII)




(wherein

R_3 is optionally alkyl-substituted 1,2-cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R_4 is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or



each R_7 independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof, said process comprising hydrogenating a compound of formula VII



(wherein R_3 , R_4 and R_7 are as hereinbefore defined) and optionally converting the compound obtained into a said salt.


14. A process for preparing a magnetic resonance imaging contrast medium composition, said process comprising admixing a chelate or a calcium salt thereof prepared according to any one of claims 7 to 11 together with a pharmaceutically acceptable, compatible excipient.

15. Use of a chelate or a calcium salt thereof as defined in any one of claims 7 to 11 for the preparation of a magnetic resonance imaging contrast medium composition.

16. A method of imaging comprising administering to a subject a chelate or a calcium salt thereof as defined in any one of claims 7 to 11 and generating an image of at least a part of said subject.


Claims for the following Contracting State : GR

1. A compound of formula I




(wherein


R is hydrogen or



R₁ is hydrogen or




with the proviso that at least one of R and R₁ is other than hydrogen;
 R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted
 1,2-cycloalkylene having from 5 to 8 carbons, or optionally alkyl substituted 1,2-arylene having from 6
 to 10 carbons; and
 R₄ is hydrogen, hydroxymethyl, alkyl having from 1 to 6 carbons or




R₅ and R₆ are each, independently, hydroxy, alkoxy having from 1 to 18 carbons, hydroxy-
 substituted alkoxy having from 1 to 18 carbons, amino or alkylamido having from 1 to carbons; and
 each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18
 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.

2. A compound as claimed in Claim 1 wherein for formula I each R₇ is hydrogen.


3. A compound as claimed in Claim 1 wherein for formula I R is



and R₁ is



4. A compound as claimed in Claim 3 wherein for formula I R₅ and R₆ are each, independently, hydroxy, alkoxy having from 1 to 8 carbons, amino or alkylamido having from 1 to 8 carbons.
5. A compound as claimed in Claim 4 wherein for formula I R₅ and R₆ are hydroxy.
6. A compound as claimed in Claim 5 being N,N'-bis-(pyridoxal-5-phosphate)ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.
7. A chelate of a chelating compound as claimed in any one of claims 1 to 6 with a metal ion having an atomic number of from 21 to 29, 42, 44 or 58-70, or a salt thereof.
8. A chelate as claimed in Claim 7 wherein the metal ion is selected from chromium(III), manganese(II), iron(III), iron(II), cobalt(II), nickel(II), copper(II), praseodymium(III), neodymium(III), samarium(III), ytterbium(III), gadolinium(III), terbium(III), dysprosium(III), holmium(III) and erbium(III).
9. A chelate as claimed in Claim 8 being a manganese(II) chelate of N,N'-bis-(pyridoxal-5-phosphate)-ethylenediamine-N,N'-diacetic acid or a salt thereof, or N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylene-diamine-N,N'-diacetic acid or a salt thereof.
10. A calcium salt of a chelate as claimed in any one of Claims 7 to 9.
11. A calcium salt as claimed in Claim 10 wherein the molar ratio of calcium to the chelating compound of formula I is from 0.05 to 1.0.
12. A chelating compound intermediate of formula (VII)




(VII)

wherein


R₃ is alkylene having from 1 to 8 carbons, optionally alkyl-substituted cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R₄ is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or



each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.

13. A chelating compound intermediate of formula (VIII)




(VIII)

wherein

R₃ is optionally alkyl-substituted 1,2-cycloalkylene having from 5 to 8 carbons or optionally alkyl-substituted 1,2-arylene having from 6 to 10 carbons;

R₄ is hydrogen, hydroxyalkyl having from 1 to 6 carbons, alkyl having from 1 to 6 carbons or



each R₇ independently is hydrogen, optionally hydroxy-substituted alkyl having from 1 to 18 carbons, optionally hydroxy-substituted alkylamino alkyl having from 1 to 18 carbons) or a salt thereof.

14. A process for preparing a magnetic resonance imaging contrast medium composition, said process comprising admixing a chelate or a calcium salt thereof as claimed in any one of claims 7 to 11 together with a pharmaceutically acceptable, compatible excipient.


15. Use of a chelate or a calcium salt thereof as defined in any one of claims 7 to 11 for the preparation of a magnetic resonance imaging contrast medium composition.

16. A method of imaging comprising administering to a subject a chelate or a calcium salt thereof as claimed in any one of claims 7 to 11 and generating an image of at least a part of said subject.


Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL, SE


1. Eine Verbindung der Formel




(worin
R Wasserstoff oder



ist;
R₁ Wasserstoff oder



bedeutet,
mit der Maßgabe, daß wenigstens eine der Gruppen R und R₁ von Wasserstoff verschieden ist;
R₃ Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen bedeutet, und
R₄ Wasserstoff, Hydroxymethyl, Alkyl mit 1 bis 6 Kohlenstoffatomen oder




bedeutet,
R₅ und R₆ unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 18 Kohlenstoffatomen, Hydroxy-substituiertes Alkoxy mit 1 bis 18 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 18 Kohlenstoffatomen bedeuten, und
jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet) oder ein Salz hiervon.

2. Eine Verbindung gemäß Anspruch 1, Worin jede Gruppe R₇ in Formel I Wasserstoff bedeutet.


3. Eine Verbindung gemäß Anspruch 1, worin in Formel I die Gruppe R

5



und R₁

10



15


bedeutet.

4. Eine Verbindung gemäß Anspruch 3, worin R₅ und R₆ in Formel I unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 8 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 8 Kohlenstoffatomen bedeuten.
- 20 5. Eine Verbindung gemäß Anspruch 4, worin R₅ und R₆ in Formel I Hydroxy bedeuten.
6. Eine Verbindung gemäß Anspruch 5, nämlich N,N'-Bis-(pyridoxal-5-phosphat)ethylendiamin-N,N'-diessigsäure oder ein Salz hiervon oder N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylen-diamin-N,N'-diessigsäure oder ein Salz hiervon.
- 25 7. Ein Chelat einer Chelatisierungsverbindung gemäß einem der Ansprüche 1 bis 6 mit einem Metallion mit einer Atomzahl von 21 bis 29, 42, 44 oder 58-70, oder ein Salz hiervon.
8. Ein Chelat gemäß Anspruch 7, worin das Metallion ausgewählt ist aus Chrom(III), Mangan(II), Eisen(III), Eisen(II), Kobalt(II), Nickel(II), Kupfer(II), Praseodym(III), Neodym(III), Samarium(III), Ytterbium(III), Gadolinium(III), Terbium(III), Dysprosium(III), Holmium(III) und Erbium(III).
- 30 9. Ein Chelat gemäß Anspruch 8, nämlich ein Mangan(II)-Chelat von N,N'-Bis-(pyridoxal-5-phosphat)-ethylendiamin-N,N'-diessigsäure oder einem Salz hiervon oder von N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylen-diamin-N,N'-diessigsäure oder einem Salz hiervon.
- 35 10. Ein Calcium-Salz eines Chelats gemäß einem der Ansprüche 7 bis 9.
11. Ein Calcium-Salz gemäß Anspruch 10, worin das molare Verhältnis von Calcium zur Chelatisierungsverbindung der Formel I 0,05 bis 1,0 beträgt.
- 40 12. Eine Chelatisierungs-Zwischenverbindung der Formel (VII)

45

50


55



worin

R₃ Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen darstellt;


R₄ Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet, oder ein Salz hiervon.

13. Eine Chelatisierungs-Zwischenverbindung der Formel (VIII)




(VIII)

worin

R₃ wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen ist;

R₄ Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen ist, oder ein Salz hiervon.

14. Eine Kontrastmittelzusammensetzung für die magnetische Resonanz-Bildgebung, umfassend einen Chelat oder ein Calcium-Salz hiervon gemäß einem der Ansprüche 7 bis 11 zusammen mit einem pharmazeutisch zulässigen, verträglichen Trägermittel.


15. Verwendung eines Chelates oder eines Calcium-Salzes hiervon, wie beschrieben in einem der Ansprüche 7 bis 11, zur Herstellung einer Kontrastmittelzusammensetzung für die magnetische Resonanz-Bildgebung.

16. Ein Verfahren zur Bildgebung, umfassend das Verabreichen eines Chelates oder eines Calcium-Salzes hiervon gemäß einem der Ansprüche 7 bis 11 an ein Subjekt und die Herstellung eines Bildes von


wenigstens einem Teil dieses Subjekts.

Patentansprüche für folgenden Vertragsstaat : ES


1. Ein Verfahren zur Herstellung einer Verbindung der Formel I



(worin
R Wasserstoff oder



ist;
R₁ Wasserstoff oder




bedeutet,

mit der Maßgabe, daß wenigstens eine der Gruppen R und R₁ von Wasserstoff verschieden ist;

R₃ Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen bedeutet, und


R₄ Wasserstoff, Hydroxymethyl, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

R₅ und R₆ unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 18 Kohlenstoffatomen, Hydroxy-substituiertes Alkoxy mit 1 bis 18 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 18 Kohlenstoffatomen bedeuten, und

jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet) oder ein Salz hiervon, umfassend die Umsetzung einer Verbindung der Formel VIII




(worin R_3 , R_4 und R_7 die oben angegebene Bedeutung haben, mit einer Halo-Essigsäure, und wahlweise die Umwandlung der erhaltenen Verbindung in ein Salz


- 15 2. Ein Verfahren gemäß Anspruch 1, worin in der Verbindung der Formel VIII jede Gruppe R_7 Wasserstoff ist.

3. Ein Verfahren gemäß Anspruch 1, worin in der Verbindung der Formel I R

20



25 und R_1



30

ist.

- 35 4. Ein Verfahren gemäß Anspruch 3, worin in der Verbindung gemäß Formel I R_5 und R_6 unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 8 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 8 Kohlenstoffatomen bedeuten.

5. Ein Verfahren gemäß Anspruch 4, worin in der Verbindung der Formel I R_5 und R_6 Hydroxy bedeuten.

- 40 6. Ein Verfahren gemäß Anspruch 5, worin die Verbindung der Formel I N,N'-Bis-(pyridoxal-5-phosphat)-ethylendiamin-N,N'-diessigsäure oder ein Salz hiervon oder N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylendiamin-N,N'-diessigsäure oder ein Salz hiervon ist.

- 45 7. Ein Verfahren zur Herstellung eines Chelates einer Chelatisierungsverbindung gemäß einem der Ansprüche 1 bis 6 oder eines Salzes hiervon, umfassend die Metallierung der Chelatisierungsverbindung oder eines Salzes hiervon mit einem Metallion mit einer Atomzahl von 21 bis 29, 42, 44 oder 58-70.


- 50 8. Ein Verfahren gemäß Anspruch 7, worin das Metallion ausgewählt wird aus Chrom(III), Mangan(II), Eisen(III), Eisen(II), Kobalt(II), Nickel(II), Kupfer(II), Praseodym(III), Neodym(III), Samarium(III), Ytterbium(III), Gadolinium(III), Terbium(III), Dysprosium(III), Holmium(III) und Erbium(III).

- 55 9. Ein Verfahren gemäß Anspruch 8 zur Herstellung eines Mangan(II)-Chelates von N,N'-Bis-(pyridoxal-5-phosphat)-ethylendiamin-N,N'-diessigsäure oder eines Salzes hiervon oder von N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylendiamin-N,N'-diessigsäure oder eines Salzes hiervon.

10. Ein Verfahren gemäß einem der Ansprüche 7 bis 9 zur Herstellung eines Calcium-Salzes eines Chelates wie angegeben in einem der Ansprüche 7 bis 9.

11. Ein Verfahren gemäß Anspruch 10, worin im Calcium-Salz das molare Verhältnis von Calcium zur Chelatisierungsverbindung gemäß Formel I 0,05 bis 1,0 beträgt.

12. Ein Verfahren zur Herstellung einer Chelatisierungs-Zwischenverbindung der Formel (VII)




(VII)

worin


R_3 Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen darstellt;

R_4 Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R_7 unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet, oder eines Salzes hiervon, umfassend die Umsetzung einer Verbindung der Formel V




(V)

worin R_4 die oben angegebene Bedeutung hat, mit einer Verbindung der Formel VI



worin R_3 die oben genannte Bedeutung hat, und wahlweise die Umwandlung der erhaltenen Verbindung in ein Salz.

13. Ein Verfahren zur Herstellung einer Chelatisierungs-Zwischenverbindung der Formel (VIII)




(VIII)

worin


R_3 wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen ist;

R_4 Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R_7 unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen ist, oder eines Salzes hiervon, umfassend die Hydrierung einer Verbindung der Formel VII



(VII)

worin R_3 , R_4 und R_7 die oben angegebene Bedeutung haben, und wahlweise die Umwandlung der erhaltenen Verbindung in ein Salz.


14. Ein Verfahren zur Herstellung einer Kontrastmittel-Zusammensetzung für die magnetische Resonanz-bildgebung, umfassend das Zusammenmischen eines Chelates oder eines Calcium-Salzes hiervon, hergestellt gemäß einem der Ansprüche 7 bis 11, mit einem pharmazeutisch zulässigen, verträglichen Trägermittel.

15. Verwendung eines Chelates oder eines Calcium-Salzes hiervon, wie beschrieben in einem der Ansprüche 7 bis 11, zur Herstellung einer Kontrastmittelzusammensetzung für die magnetische Resonanz-Bildgebung.


16. Ein Verfahren zur Bildgebung, umfassend das Verabreichen eines Chelates oder eines Calcium-Salzes hiervon gemäß einem der Ansprüche 7 bis 11 an ein Subjekt und die Herstellung eines Bildes von wenigstens einem Teil dieses Subjekts.

5 Patentansprüche für folgenden Vertragsstaat : GR

1. Eine Verbindung der Formel



(worin
25 R Wasserstoff oder



ist;
R₁ Wasserstoff oder



40 bedeutet,
mit der Maßgabe, daß wenigstens eine der Gruppen R und R₁ von Wasserstoff verschieden ist;
R₃ Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen bedeutet, und


45 R₄ Wasserstoff, Hydroxymethyl, Alkyl mit 1 bis 6 Kohlenstoffatomen oder




bedeutet,
R₅ und R₆ unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 18 Kohlenstoffatomen, Hydroxy-substituiertes Alkoxy mit 1 bis 18 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 18 Kohlenstoffatomen bedeuten, und
55 jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet) oder ein Salz hiervon.

2. Eine Verbindung gemäß Anspruch 1, worin jede Gruppe R_7 in Formel I Wasserstoff bedeutet.

3. Eine Verbindung gemäß Anspruch 1, worin in Formel I die Gruppe R



und R_1



bedeutet.

4. Eine Verbindung gemäß Anspruch 3, worin R_5 und R_6 in Formel I unabhängig voneinander Hydroxy, Alkoxy mit 1 bis 8 Kohlenstoffatomen, Amino oder Alkylamido mit 1 bis 8 Kohlenstoffatomen bedeuten.

5. Eine Verbindung gemäß Anspruch 4, worin R_5 und R_6 in Formel I Hydroxy bedeuten.

6. Eine Verbindung gemäß Anspruch 5, nämlich N,N'-Bis-(pyridoxal-5-phosphat)ethylendiamin-N,N'-diessigsäure oder ein Salz hiervon oder N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylen-diamin-N,N'-diessigsäure oder ein Salz hiervon.

7. Ein Chelat einer Chelatisierungsverbindung gemäß einem der Ansprüche 1 bis 6 mit einem Metallion mit einer Atomzahl von 21 bis 29, 42, 44 oder 58-70, oder ein Salz hiervon.


8. Ein Chelat gemäß Anspruch 7, worin das Metallion ausgewählt ist aus Chrom(III), Mangan(II), Eisen(III), Eisen(II), Kobalt(II), Nickel(II), Kupfer(II), Praseodym(III), Neodym(III), Samarium(III), Ytterbium(III), Gadolinium(III), Terbium(III), Dysprosium(III), Holmium(III) und Erbium(III).

9. Ein Chelat gemäß Anspruch 8, nämlich ein Mangan(II)-Chelat von N,N'-Bis-(pyridoxal-5-phosphat)-ethylendiamin-N,N'-diessigsäure oder einem Salz hiervon oder von N,N'-Bis-(pyridoxal-5-phosphat)-trans-1,2-cyclohexylen-diamin-N,N'-diessigsäure oder einem Salz hiervon.

10. Ein Calcium-Salz eines Chelats gemäß einem der Ansprüche 7 bis 9.

11. Ein Calcium-Salz gemäß Anspruch 10, worin das molare Verhältnis von Calcium zur Chelatisierungsverbindung der Formel I 0,05 bis 1,0 beträgt.


12. Eine Chelatisierungs-Zwischenverbindung der Formel (VII)



worin

R₃ Alkylen mit 1 bis 8 Kohlenstoffatomen, wahlweise Alkyl-substituiertes Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen darstellt;


R₄ Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen bedeutet, oder ein Salz hiervon.

13. Eine Chelatisierungs-Zwischenverbindung der Formel (VIII)




(VIII)

worin

R₃ wahlweise Alkyl-substituiertes 1,2-Cycloalkylen mit 5 bis 8 Kohlenstoffatomen oder wahlweise Alkyl-substituiertes 1,2-Arylen mit 6 bis 10 Kohlenstoffatomen ist;

R₄ Wasserstoff, Hydroxyalkyl mit 1 bis 6 Kohlenstoffatomen, Alkyl mit 1 bis 6 Kohlenstoffatomen oder



bedeutet,

jede Gruppe R₇ unabhängig voneinander Wasserstoff, wahlweise Hydroxy-substituiertes Alkyl mit 1 bis 18 Kohlenstoffatomen, wahlweise Hydroxy-substituiertes Alkylamino-alkyl mit 1 bis 18 Kohlenstoffatomen ist, oder ein Salz hiervon.

14. Ein Verfahren zur Herstellung einer Kontrastmittel-Zusammensetzung für die magnetische Resonanz-bildgebung, umfassend das Zusammenmischen eines Chelates oder eines Calcium-Salzes hiervon, hergestellt gemäß einem der Ansprüche 7 bis 11, mit einem pharmazeutisch zulässigen, verträglichen Trägermittel.


15. Verwendung eines Chelates oder eines Calcium-Salzes hiervon, wie beschrieben in einem der Ansprüche 7 bis 11, zur Herstellung einer Kontrastmittelzusammensetzung für die magnetische Resonanz-Bildgebung.

16. Ein Verfahren zur Bildgebung, umfassend das Verabreichen eines Chelates oder eines Calcium-Salzes hiervon gemäß einem der Ansprüche 7 bis 11 an ein Subjekt und die Herstellung eines Bildes von wenigstens einem Teil dieses Subjekts.


5 Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL, SE


1. Composé de formule I



(dans laquelle
R est un atome d'hydrogène ou




R₁ est un atome d'hydrogène ou



sous réserve qu'au moins un de R et de R₁ soit autre qu'un atome d'hydrogène ;

R₃ est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes ; et

R₄ est un atome d'hydrogène, un groupe hydroxyméthyle, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



R₅ et R₆ sont chacun indépendamment un groupe hydroxy, un groupe alcoxy ayant de 1 à 18 atomes de carbone, un groupe alcoxy hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe amino ou un groupe alkylamido ayant de 1 à 18 atomes de carbone ; et

chaque R₇, indépendamment, est un atome d'hydrogène, un groupe alkyle facultativement hy-

droxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle ayant de 1 à 18 atomes de carbone facultativement hydroxy-substitué)
ou un de ses sels.

5 2. Composé selon la revendication 1, où, dans la formule I, chaque R₇ est un atome d'hydrogène.

3. Composé selon la revendication 1, où, dans la formule I, R est



15 et R₁ est



25 4. Composé selon la revendication 3, où, dans la formule I, R₅ et R₆ sont chacun, indépendamment, un groupe hydroxy, un groupe alcoxy ayant de 1 à 8 atomes de carbone, un groupe amino ou un groupe alkylamido ayant de 1 à 8 atomes de carbone.

5. Composé selon la revendication 4, où, dans la formule I, R₅ et R₆ sont un groupe hydroxy.

30 6. Composé selon la revendication 5 qui est l'acide N,N'-bis-(pyridoxal-5-phosphate)éthylènediamine-N,N'-diacétique ou un de ses sels ou l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou un de ses sels.

35 7. Chélate d'un composé chélatant selon l'une quelconque des revendications 1 à 6 avec un ion métallique ayant un numéro atomique de 21 à 29, 42, 44 ou 58 à 70 ou un de ses sels.

8. Chélate selon la revendication 7, dans lequel l'ion métallique est choisi parmi le chrome(III), le manganèse(II), le fer(III), le fer(II), le cobalt(II), le nickel(II), le cuivre(II), le praséodyme(III), le néodyme(III), le samarium(III), l'ytterbium(III), le gadolinium(III), le terbium(III), le dysprosium(III), l'holmium(III) et l'erbium(III).


9. Chélate selon la revendication 8, qui est un chélate de manganèse(II) de l'acide N,N'-bis-(pyridoxal-5-phosphate)éthylènediamine-N,N'-diacétique ou d'un de ses sels ou de l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou d'un de ses sels.

10. Sel de calcium d'un chélate selon l'une quelconque des revendications 7 à 9.

50 11. Sel de calcium selon la revendication 10, dans lequel le rapport molaire du calcium ou composé chélatant de formule I est de 0,05 à 1,0.

55


12. Intermédiaire de composé chélatant de formule (VII)



15 (dans laquelle


R_3 est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;

R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone) ou un de ses sels.


13. Intermédiaire de composé chélatant de formule (VIII)



45 (dans laquelle

R_3 est un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;

R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone) ou un de ses sels.


14. Composition de milieu de contraste pour l'imagerie par résonance magnétique comprenant un chélate ou un sel de calcium de celui-ci selon l'une quelconque des revendications 7 à 11 avec un excipient compatible pharmaceutiquement acceptable.

15. Utilisation d'un chélate ou d'un de ses sels de calcium comme défini dans l'une quelconque des revendications 7 à 11 pour la préparation d'une composition de milieu de contraste pour l'imagerie par résonance magnétique.


16. Procédé d'imagerie comprenant l'administration à un sujet d'un chélate ou d'un de ses sels de calcium selon l'une quelconque des revendications 7 à 11 et l'établissement d'une image d'au moins une partie dudit sujet.

Revendications pour l'Etat contractant suivant : ES


1. Procédé pour préparer un composé de formule I



(dans laquelle
R est un atome d'hydrogène ou




R_1 est un atome d'hydrogène ou



sous réserve qu'au moins un de R et de R₁ soit autre qu'un atome d'hydrogène ;

R₃ est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ; et

R₄ est un atome d'hydrogène, un groupe hydroxyméthyle, un groupe alkyle ayant de 1 à 6 atomes de carbone ou




R₅ et R₆ sont chacun indépendamment un groupe hydroxy, un groupe alcoxy ayant de 1 à 18 atomes de carbone, un groupe alcoxy hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe amino ou un groupe alkylamido ayant de 1 à 18 atomes de carbone ; et

chaque R₇, indépendamment, est un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone)

ou un de ses sels,


ledit procédé comprenant la réaction d'un composé de formule VIII




(dans laquelle R₃, R₄ et R₇ sont tels que précédemment définis) avec un acide halogénoacétique et facultativement la conversion du composé obtenu en ledit sel.

2. Procédé selon la revendication 1, où, dans ledit composé de formule VIII, chaque R₇ est un atome d'hydrogène.

3. Procédé selon la revendication 1, où, dans ledit composé de formule I, R est




et R₁ est



4. Procédé selon la revendication 3, où, dans ledit composé de formule I, R₅ et R₆ sont chacun, indépendamment, un groupe hydroxy, un groupe alcoxy ayant de 1 à 8 atomes de carbone, un groupe

amino ou un groupe alkylamido ayant de 1 à 8 atomes de carbone.

5. Procédé selon la revendication 4, où, dans ledit composé de formule I, R_5 et R_6 sont un groupe hydroxy.
6. Procédé selon la revendication 5 où ledit composé de formule I est l'acide N,N'-bis-(pyridoxal-5-phosphate)éthylènediamine-N,N'-diacétique ou un de ses sels ou l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou un de ses sels.
7. Procédé pour préparer un chélate d'un composé chélatant selon l'une quelconque des revendications 1 à 6 ou d'un sel de celui-ci, ledit procédé comprenant la métallation dudit composé chélatant ou d'un de ses sels avec un ion métallique ayant un numéro atomique de 21 à 29, 42, 44 ou 58 à 70.
8. Procédé selon la revendication 7, dans lequel l'ion métallique est choisi parmi le chrome(III), le manganèse(II), le fer(III), le fer(II), le cobalt(II), le nickel(II), le cuivre(II), le praséodyme(III), le néodyme(III), le samarium(III), l'ytterbium(III), le gadolinium(III), le terbium(III), le dysprosium(III), l'holmium(III) et l'erbium(III).
9. Procédé selon la revendication 8 pour préparer un chélate de manganèse(II) de l'acide N,N'-bis-(pyridoxal-5-phosphate)éthylènediamine-N,N'-diacétique ou d'un de ses sels ou de l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou d'un de ses sels.
10. Procédé selon l'une quelconque des revendications 7 à 9 pour préparer un sel de calcium d'un chélate tel que défini dans l'une quelconque des revendications 7 à 9.
11. Procédé selon la revendication 10, où, dans ledit sel de calcium, le rapport molaire du calcium au composé chélatant de formule I est de 0,05 à 1,0.
12. Procédé pour préparer un intermédiaire de composé chélatant de formule (VII)




(VII)

(dans laquelle


R_3 est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;

R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement

hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone) ou un de ses sels, ledit procédé comprenant la réaction d'un composé de formule V




(V)

(dans laquelle R_4 est tel que précédemment défini) avec un composé de formule VI



(dans laquelle R_3 est tel que précédemment défini) et facultativement la conversion du composé obtenu en ledit sel.

13. Procédé pur préparer un intermédiaire de composé chélatant de formule (VIII)




(VIII)

(dans laquelle

R_3 est un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;


R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone)

ou un de ses sels,

ledit procédé comprenant l'hydrogénation d'un composé de formule (VII)




(VII)

(dans laquelle R_3 , R_4 et R_7 sont tels que précédemment définis) et facultativement la conversion du composé obtenu en ledit sel.

14. Procédé pour préparer une composition de milieu de contraste pour l'imagerie par résonance magnétique, ledit procédé comprenant le mélange d'un chélate, ou d'un sel de calcium de celui-ci, préparé selon l'une quelconque des revendications 7 à 11 avec un excipient compatible pharmaceutiquement acceptable.
15. Utilisation d'un chélate ou d'un de ses sels de calcium comme défini dans l'une quelconque des revendications 7 à 11 pour la préparation d'une composition de milieu de contraste pour l'imagerie par résonance magnétique.
16. Procédé d'imagerie comprenant l'administration à un sujet d'un chélate ou d'un de ses sels de calcium comme défini dans l'une quelconque des revendications 7 à 11 et l'établissement d'une image d'au moins une partie dudit sujet.


Revendications pour l'Etat contractant suivant : GR

1. Composé de formule I




(I)

(dans laquelle
R est un atome d'hydrogène ou



R_1 est un atome d'hydrogène ou




5

sous réserve qu'au moins un de R et de R₁ soit autre qu'un atome d'hydrogène ;

R₃ est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ; et

10

R₄ est un atome d'hydrogène, un groupe hydroxyméthyle, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



15

R₅ et R₆ sont chacun indépendamment un groupe hydroxy, un groupe alcoxy ayant de 1 à 18 atomes de carbone, un groupe alcoxy hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe amino ou un groupe alkylamido ayant de 1 à 18 atomes de carbone ; et


20

chaque R₇, indépendamment, est un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle ayant de 1 à 18 atomes de carbone facultativement hydroxy-substitué) ou un de ses sels.

25


2. Composé selon la revendication 1, où, dans la formule I, chaque R₇ est un atome d'hydrogène.

30 3. Composé selon la revendication 1, où, dans la formule I, R est



35

et R₁ est



40

45

4. Composé selon la revendication 3, où, dans la formule I, R₅ et R₆ sont chacun, indépendamment, un groupe hydroxy, un groupe alcoxy ayant de 1 à 8 atomes de carbone, un groupe amino ou un groupe alkylamido ayant de 1 à 8 atomes de carbone.

50

5. Composé selon la revendication 4, où, dans la formule I, R₅ et R₆ sont un groupe hydroxy.

6. Composé selon la revendication 5 qui est l'acide N,N'-bis-(pyridoxal-5-phosphate)-éthylènediamine-N,N'-diacétique ou un de ses sels ou l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou un de ses sels.

55

7. Chélate d'un composé chélatant selon l'une quelconque des revendications 1 à 6 avec un ion métallique ayant un numéro atomique de 21 à 29, 42, 44 ou 58 à 70 ou un de ses sels.


8. Chélate selon la revendication 7, dans lequel l'ion métallique est choisi parmi le chrome(III), le manganèse(II), le fer(III), le fer(II), le cobalt(II), le nickel(II), le cuivre(II), le praséodyme(III), le néodyme(III), le samarium(III), l'ytterbium(III), le gadolinium(III), le terbium(III), le dysprosium(III), l'holmium(III) et l'erbium(III).

9. Chélate selon la revendication 8, qui est un chélate de manganèse(II) de l'acide N,N'-bis-(pyridoxal-5-phosphate)éthylènediamine-N,N'-diacétique ou d'un de ses sels ou de l'acide N,N'-bis-(pyridoxal-5-phosphate)-trans-1,2-cyclohexylènediamine-N,N'-diacétique ou d'un de ses sels.

10. Sel de calcium d'un chélate selon l'une quelconque des revendications 7 à 9.

11. Sel de calcium selon la revendication 10, dans lequel le rapport molaire du calcium au composé chélatant de formule I est de 0,05 à 1,0.

12. Intermédiaire de composé chélatant de formule (VII)




(VII)

(dans laquelle


R_3 est un groupe alkylène ayant de 1 à 8 atomes de carbone, un groupe cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;

R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone) ou un de ses sels.

13. Intermédiaire de composé chélatant de formule (VIII)




(VIII)

(dans laquelle

R_3 est un groupe 1,2-cycloalkylène facultativement alkyl-substitué ayant de 5 à 8 atomes de carbone ou un groupe 1,2-arylène facultativement alkyl-substitué ayant de 6 à 10 atomes de carbone ;

R_4 est un atome d'hydrogène, un groupe hydroxyalkyle ayant de 1 à 6 atomes de carbone, un groupe alkyle ayant de 1 à 6 atomes de carbone ou



chaque R_7 est, indépendamment, un atome d'hydrogène, un groupe alkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone, un groupe alkylaminoalkyle facultativement hydroxy-substitué ayant de 1 à 18 atomes de carbone) ou un de ses sels.

14. Procédé pour préparer une composition de milieu de contraste pour l'imagerie par résonance magnétique, ledit procédé comprenant le mélange d'un chélate ou d'un sel de calcium de celui-ci selon l'une quelconque des revendications 7 à 11 avec un excipient compatible pharmaceutiquement acceptable.

15. Utilisation d'un chélate ou d'un de ses sels de calcium comme défini dans l'une quelconque des revendications 7 à 11 pour la préparation d'une composition de milieu de contraste pour l'imagerie par résonance magnétique.

16. Procédé d'imagerie comprenant l'administration à un sujet d'un chélate ou d'un de ses sels de calcium selon l'une quelconque des revendications 7 à 11 et l'établissement d'une image d'au moins une partie dudit sujet.




FIG. 1.




FIG. 2.